Intiracetam Concentrate for Solution for Infusion 100mg/ml

PHARMACEUTICAL FORM AND STRENGTH

Levetiracetam, 100mg/ml, concentrate for solution for infusion Clear and colourless solution free from visible particulate matter.

EXCIPIENTS

Sodium acetate trihydrate, Sodium chloride, Glacial acetic acid, Water for injection

INDICATIONS

Intiracetam is indicated as monotherapy in the treatment of partial onset seizures with or without secondary generalisation in adults and adolescents from 16 years of age with newly diagnosed epilepsy.

Intiracetam is indicated as adjunctive therapy in the treatment of:

- partial onset seizures with or without secondary generalisation in adults, adolescents and children from 4 years of age with epilepsy
- myoclonic seizures in adults and adolescents from 12 years of age with Juvenile Myoclonic Epilepsy
- primary generalised tonic-clonic seizures in adults and adolescents from 12 years of age with Idiopathic Generalised Epilepsy

Intiracetam Concentrate is an alternative for patients (adults and children from 4 years of age) when oral administration is temporarily not feasible.

DOSAGE AND ADMINISTRATION

Intiracetam therapy can be initiated with either intravenous or oral administration. Conversion to or from oral to intravenous administration can be done directly without titration. The total daily dose and frequency of administration should be maintained.

Concentrate for solution for infusion

Intiracetam concentrate is for intravenous use only and the recommended dose must be diluted in at least 100 ml of a compatible diluent and administered intravenously as 15-minute intravenous infusion (*see Section Incompatibilities and Use and Handling*). There is no experience with administration of intravenous Intiracetam for longer period than 4 days.

Intiracetam concentrate is an alternative for patients (adults and children from 4 years of age) when oral administration is temporarily not feasible.

Please refer to Section INCOMPATIBILITIES AND USE AND HANDLING for recommendation on the preparation and administration of Intiracetam Concentrate for Solution for Infusion 100mg/ml.

Adults

- Monotherapy

  Adults and adolescents from 16 years of age

  The recommended starting dose is 250 mg twice daily which should be increased to an initial therapeutic dose of 500 mg twice daily after two weeks. The dose can be further increased by 250 mg twice daily every two weeks depending upon the clinical response. The maximum dose is 1500 mg twice daily.

  The safety and efficacy of levetiracetam in children and adolescents below 16 years as monotherapy

treatment have not been established. There are no data available.

- Add-on therapy

  Adults (≥18 years) and adolescents (12 to 17 years) weighing 50 kg or more

  The initial therapeutic dose is 500mg twice daily. This dose can be started on the first day of treatment. Depending upon the clinical response and tolerance, the daily dose can be increased up to 1,500mg twice daily. Dose changes can be made in 500mg twice daily increases or decreases every two to four weeks.

Children

The safety and efficacy of Intiracetam Concentrate for Solution for Infusion in infants and children less than 4 years have not been established.

- Monotherapy

  The safety and efficacy of levetiracetam in children and adolescents below 16 years as monotherapy treatment have not been established.

  There are no data available.

- Add-on therapy for children (4 to 11 years) and adolescents (12 to 17 years) weighing less than 50kg:

  The initial therapeutic dose is 10 mg/kg twice daily.

  Depending upon the clinical response and tolerability, the dose can be increased up to 30 mg/kg twice daily. Dose changes should not exceed increases or decreases of 10 mg/kg twice daily every two weeks. The lowest effective dose should be used. Dose in children 50 kg or greater is the same as in adults.

Dose recommendations for children and adolescents:

Weight	Starting dose: 10 mg/kg twice daily	Maximum dose: 30 mg/kg twice daily
15 kg(1)	150 mg (1.5ml) twice daily	450 mg (4.5ml) twice daily
20 kg(1)	200 mg(2ml) twice daily	600 mg (6ml) twice daily
25 kg	250 mg twice daily	750 mg twice daily
From 50 kg(2)	500 mg twice daily	1500 mg twice daily

(1) Children 25kg or less should preferably start the treatment with levetiracetam 100mg/ml oral solution

(2) Dosage in children and adolescents 50 kg or more is the same as in adults.

Infants and children less than 4 years

There are insufficient data to recommend the use of levetiracetam in children under 4 years of age.

For children with renal impairment, Intiracetam dose needs to be adjusted based on the renal function as Intiracetam clearance is related to renal function. This recommendation is based on a study in adult renally impaired patients.

Elderly

Adjustment of the dose is recommended in elderly patients with compromised renal function.

Renal impairment

The daily dose must be individualised according to renal function (see section Warnings and Precautions). For adult patients, refer to the following table and adjust the dose as indicated. To use this dosing table, an estimate of the patient's creatinine clearance (CLcr) in ml/min is needed. The CLcr in ml/min may be estimated from serum creatinine (mg/dl) determination, for adults and adolescents weighing 50 kg or more, using the following formula:

$$CL_{cr} = \frac{[140\text{-age (years)}] \times \text{weight (kg)}}{72 \times \text{serum creatinine (mg/dl)}} \text{ (x 0.85 for women)}$$

Then CL_{cr} is adjusted for body surface area (BSA) as follows:

$$CL_{cr}\,(ml/min/1.73m^2) = \frac{CL_{cr}\,(ml/min)}{\text{BSA subject } (m^2)} \times 1.73$$

Dosing adjustment for adult and adolescent patients weighing more than 50 kg with impaired renal function

Group	Creatinine clearance (ml/min/1.73m^2)	Dosage and frequency
Normal	> 80	500 to 1500 mg twice daily
Mild	50-79	500 to 1000 mg twice daily
Moderate	30-49	250 to 750 mg twice daily
Severe	< 30	250 to 500 mg twice daily
End-stage renal disease patients undergoing dialysis[1]	-	500 to 1000 mg once daily[2]

[1] A 750 mg loading dose is recommended on the first day of treatment with Intiracetam.

[2] Following dialysis, a 250 to 500 mg supplemental dose is recommended.

Hepatic impairment

No dose adjustment is needed in patients with mild to moderate hepatic impairment. In patients with severe hepatic impairment, the creatinine clearance may underestimate the renal insufficiency. Therefore a 50% reduction of the daily maintenance dose is recommended when the creatinine clearance is <60 ml/min/1.73m^2

CONTRAINDICATIONS

Hypersensitivity to the active substance or other pyrrolidone derivatives or to any of the excipients.

WARNINGS AND PRECAUTIONS

Discontinuation

If Intiracetam has to be discontinued, it is recommended to withdraw it gradually (e.g. in adults and adolescents weighing 50kg or more: reduce dose by 500mg twice daily every two to four weeks: children and adolescents weighing less than 50kg: dose reductions should not exceed 10 mg/kg twice daily every two weeks).

An increase in seizure frequency of more than 25% has been reported in 14% of levetiracetam treated adult and paediatric patients with partial onset seizures, whereas it was reported in 26% and 21% of placebo treated adult and paediatric patients, respectively.

When Intiracetam was used to treat primary tonic-clonic seizures in adults and adolescents with idiopathic generalised epilepsy, there was no effect on the frequency of absences.

Renal or hepatic impairment

The administration of Intiracetam to patients with renal impairment may require dose adjustment. In patients with severely impaired hepatic function, assessment of renal function is recommended before dose selection *(see section Dosage and Administration)*.

Acute Kidney Injury

The use of Intiracetam has been rarely associated with acute kidney injury, with a time to onset ranging

from a few days to several months.

Blood cell counts

Rare cases of decreased blood cell counts (neutropenia, agranulocytosis, leucopenia, thrombocytopenia and pancytopenia) have been described in association with Intiracetam administration, generally at the beginning of the treatment. Complete blood cell counts are advised in patients experiencing important weakness, pyrexia, recurrent infections or coagulation disorders (*see Section Adverse Reactions)*.

Psychiatric Reactions

Behavioural abnormalities including psychotic symptoms, suicidal ideation, irritability and aggressive behaviour have been observed. Monitor patients for psychiatric signs and symptoms. If such behaviours are noticed, treatment adaptation or gradual discontinuation should be considered. If discontinuation is considered, please see Section Discontinuation in Warnings and Precautions.

A total of 13.3% of adult levetiracetam-treated patients and 37.6% of paediatric levetiracetam-treated patients (4 to 16 years of age) compared to 6.2% and 18.6% of adult and paediatric placebo patients respectively, experienced non-psychotic behavioural symptoms (reported as aggression, agitation, anger, anxiety, apathy, depersonalization, depression, emotional lability, hostility, hyperkinesias, irritability, nervousness, neurosis, and personality disorder).

A total of 1.7% of adult levetiracetam-treated patients discontinued treatment due to behavioural adverse events, compared to 0.2% of placebo patients. The treatment dose was reduced in 0.8% of adult levetiracetam-treated patients and in 0.5% of placebo patients. Overall, 10.9% of levetiracetam-treated pediatric patients experiences behavioural symptoms associated with discontinuation or dose reduction, compared to 6.2% of placebo patients.

One percent of adult levetiracetam-treated patients experienced psychotic symptoms compared to 0.2% in the placebo patients.

Two (0.3%) adult levetiracetam-treated patients were hospitalized and their treatment was discontinued due to psychosis. Both events, reported as psychosis, developed within the first week of treatment and resolved within 1 to 2 weeks following treatment discontinuation. There was no difference between drug and placebo-treated patients in the incidence of the paediatric patients who discontinued treatment due to psychotic and non-psychotic adverse reactions.

Depression and/ or suicidal ideation

Suicide, suicide attempt and suicidal ideation and behaviour have been reported in patients treated with anti-epileptic agents (including levetiracetam). A meta-analysis of randomised placebo-controlled trials of anti-epileptic medicinal products has shown a small increased risk of suicidal thoughts and behaviour. The mechanism of this risk is not known. Therefore patients should be monitored for signs of depression and/or suicidal ideation and behaviours and appropriate treatment should be considered. Patients (and caregivers of patients) should be advised to seek medical advice should signs of depression and/or suicidal ideation or behaviour emerge.

Paediatric population

Available data in children did not suggest impact on growth and puberty. However, long-term effects on learning, intelligence, growth, endocrine function, puberty and child bearing potential in children remain unknown.

Excipients

This medicinal product contains 2.5 mmol (or 57 mg) sodium per maximum single dose (0.8 mmol (or 19 mg) per vial). To be taken into consideration by patients on a controlled sodium diet.

INTERACTIONS

Anti-epileptic medicinal products

Pre-marketing data from clinical studies conducted in adults indicate that levetiracetam did not influence the serum concentrations of existing anti-epileptic medicinal products (phenytoin, carbamazepine, valproic acid, phenobarbital, lamotrigine, gabapentin and primidone) and that these anti-epileptic medicinal products did not influence the pharmacokinetics of levetiracetam.

As in adults, there is no evidence of clinically significant medicinal product interactions in paediatric patients receiving up to 60 mg/kg/day levetiracetam.

A retrospective assessment of pharmacokinetic interactions in children and adolescents with epilepsy (4 to 17 years) confirmed that adjunctive therapy with levetiracetam did not influence the steady-state serum concentrations of concomitantly administered carbamazepine and valproate. However, data suggested a 20% higher levetiracetam clearance in children taking enzyme-inducing anti-epileptic medicinal products. Dose adjustment is not required.

Probenecid

Probenecid (500 mg four times daily), a renal tubular secretion blocking agent, has been shown to inhibit the renal clearance of the primary metabolite but not of levetiracetam. Nevertheless, the concentration of this metabolite remains low.

Methotrexate

Concomitant administration of levetiracetam and methotrexate has been reported to decrease methotrexate clearance, resulting in increased/prolonged blood methotrexate concentration to potentially toxic levels. Blood methotrexate and levetiracetam levels should be carefully monitored in patients treated concomitantly with the two drugs.

Oral contraceptives and other pharmacokinetics interactions

Levetiracetam 1000 mg daily did not influence the pharmacokinetics of oral contraceptives (ethinyl- estradiol and levonorgestrel); endocrine parameters (luteinising hormone and progesterone) were not modified. Levetiracetam 2000 mg daily did not influence the pharmacokinetics of digoxin and warfarin; prothrombin times were not modified. Co-administration with digoxin, oral contraceptives and warfarin did not influence the pharmacokinetics of levetiracetam.

Alcohol

No data on the interaction of levetiracetam with alcohol are available.

PREGNANCY AND LACTATION

Women of child bearing potential

Specialist advice should be given to women who are of childbearing potential. Treatment with levetiracetam should be reviewed when a woman is planning to become pregnant. As with all antiepileptic medicines, sudden discontinuation of levetiracetam should be avoided as this may lead to breakthrough seizures that could have serious consequences for the woman and the unborn child. Monotherapy should be preferred whenever possible because therapy with multiple antiepileptic medicines AEDs could be associated with a higher risk of congenital malformations than monotherapy, depending on the associated antiepileptics.

Pregnancy

A large amount of postmarketing data on pregnant women exposed to levetiracetam monotherapy (more than 1800, among which in more than 1500 exposure occurred during the 1st trimester) do not suggest an increase in the risk for major congenital malformations. Only limited evidence is available on the neurodevelopment of children exposed to levetiracetam monotherapy in utero. However, current epidemiological studies (on about 100 children) do not suggest an increased risk of neurodevelopmental disorders or delays. Studies in animals have shown reproductive toxicity.

Levetiracetam can be used during pregnancy, if after careful assessment it is considered clinically needed. In such case, the lowest effective dose is recommended.

Physiological changes during pregnancy may affect levetiracetam concentration. Decrease in levetiracetam plasma concentrations has been observed during pregnancy. This decrease is more pronounced during the third trimester (up to 60% of baseline concentration before pregnancy). Appropriate clinical management of pregnant women treated with levetiracetam should be ensured.

Breastfeeding

Levetiracetam is excreted in human breast milk. Therefore, breast-feeding is not recommended.

However, if levetiracetam treatment is needed during breastfeeding, the benefit/risk of the treatment should be weighed considering the importance of breastfeeding.

Fertility

No impact on fertility was detected in animal studies. No clinical data are available, potential risk for human is unknown.

ABILITY TO PERFORM TASKS THAT REQUIRE JUDGEMENT, MOTOR OR COGNITIVE SKILLS

Levetiracetam has minor or moderate influence on the ability to drive and use machines. Due to possible different individual sensitivity, some patients might experience somnolence or other central nervous system related symptoms, at the beginning of treatment or following a dose increase. Therefore, caution is recommended in those patients when performing skilled tasks, e.g. driving vehicles or operating machinery. Patients are advised not to drive or use machines until it is established that their ability to perform such activities is not affected.

ADVERSE REACTIONS

Clinical Trial Data and Post-Marketing Data

Summary of the safety profile

The adverse event profile presented below is based on the analysis of pooled placebo-controlled clinical trials with all indications studied, with a total of 3,416 patients treated with levetiracetam. These data are supplemented with the use of levetiracetam in corresponding open-label extension studies, as well as post-marketing experience.

The most frequently reported adverse reactions were nasopharyngitis, somnolence, headache, fatigue and dizziness. The safety profile of levetiracetam is generally similar across age groups (adult and paediatric patients) and across the approved epilepsy indications.

Tabulated list of adverse reactions

Adverse reactions are presented in the order of decreasing seriousness and their frequency is defined as follows: very common (≥1/10); common (≥1/100 to <1/10); uncommon (≥1/1,000 to <1/100); rare (≥ 1/10,000 to <1/1,000) and very rare (<1/10,000).

MedDRA SOC	Frequency category			
	Very common	Common	Uncommon	Rare
Infections and infestations	Nasopharyngitis			Infection
Blood and lymphatic system disorders			Thrombocytopenia, leukopenia	Pancytopenia, neutropenia, agranulocytosis
Immune system disorders				Drug reaction with eosinophilia and systemic symptoms (DRESS), Hypersensitivity (including angioedema and anaphylaxis)
Metabolism and nutrition disorders		Anorexia	Weight decreased, weight increase	Hyponatraemia
Psychiatric disorders		Depression, hostility/ aggression, anxiety, insomnia, nervousness/irritability	Suicide attempt, suicidal ideation, psychotic disorder, abnormal behaviour, hallucination, anger, confusional state, panic attack, affect lability/mood swings,	Completed suicide, personality disorder, thinking abnormal, delirium

			agitation	
Nervous system disorders	Somnolence, headache	Convulsion, balance disorder, dizziness, lethargy, tremor	Amnesia, memory impairment, coordination abnormal/ataxia, paraesthesia, disturbance in attention	Choreoathetosis, dyskinesia, hyperkinesia, gait disturbance, encephalopathy
Eye disorders			Diplopia, vision blurred	
Ear and labyrinth disorders		Vertigo		
Respiratory, thoracic and mediastinal disorders		Cough		
Gastrointestinal disorders		Abdominal pain, diarrhoea, dyspepsia, vomiting, nausea		Pancreatitis
Hepatobiliary disorders			Liver function test abnormal	Hepatic failure, hepatitis
Renal and Urinary Disorders				Acute Kidney injury
Skin and subcutaneous tissue disorders		Rash	Alopecia, eczema, pruritus,	Toxic epidermal necrolysis, Stevens-Johnson syndrome, erythema multiforme
Musculoskeletal and connective tissue disorders			Muscular weakness, myalgia	Rhabdomyolysis and blood creatine phosphokinase increased*
General disorders and administration site conditions		Asthenia/fatigue		
Injury, poisoning and procedural complications			Injury	

* Prevalence is significantly higher in Japanese patients when compared to non-Japanese patients.

Description of selected adverse reactions

The risk of anorexia is higher when levetiracetam is coadministered with topiramate.

In several cases of alopecia, recovery was observed when levetiracetam was discontinued.

Bone marrow suppression was identified in some of the cases of pancytopenia.

Cases of encephalopathy have been rarely observed after levetiracetam administration. These undesirable effects generally occurred at the beginning of the treatment (few days to a few months) and were reversible after treatment discontinuation.

Paediatric population

In patients aged 4-16 years, a total of 645 patients have been treated with levetiracetam in placebo- controlled and open label extension studies. 233 of these patients were treated with levetiracetam in placebo-controlled studies. In this paediatric age range, the data are supplemented with post-marketing experience of the use of levetiracetam.

A double-blind, placebo-controlled paediatric safety study with a non-inferiority design has assessed the cognitive and neuropsychological effects of levetiracetam in children 4 to 16 years of age with partial onset seizures. It was concluded that levetiracetam was not different (non inferior) from placebo with regard to the change from baseline of the Leiter-R Attention and Memory, Memory Screen Composite score in the per-protocol population. Results related to behavioural and emotional functioning indicated a worsening in levetiracetam treated patients on aggressive behaviour as measured in a standardised and systematic way using

a validated instrument (CBCL – Achenbach Child Behavior Checklist). However, subjects who took levetiracetam in the long-term open label follow-up study, did not experience a worsening, on average, in their behavioural and emotional functioning; in particular measures of aggressive behaviour were not worse than baseline.

The adverse event profile of levetiracetam is generally similar across age groups and across the approved epilepsy indications. Safety results in paediatric patients in placebo-controlled clinical studies were consistent with the safety profile of levetiracetam in adults except for behavioural and psychiatric adverse reactions which were more common in children than in adults. In children and adolescents aged 4 to 16 years, vomiting (very common, 11.2%), agitation (common, 3.4%), mood swings (common, 2.1%), affect lability (common, 1.7%), aggression (common, 8.2%), abnormal behaviour (common, 5.6%), and lethargy (common, 3.9%) were reported more frequently than in other age ranges or in the overall safety profile.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system.

OVERDOSAGE

Symptoms and signs

Somnolence, agitation, aggression, depressed level of consciousness, respiratory depression and coma were observed with Intiracetam overdoses.

Management of overdose

There is no specific antidote for levetiracetam. Treatment of an overdose will be symptomatic and may include haemodialysis. The dialyser extraction efficiency is 60% for levetiracetam and 74% for the primary metabolite.

CLINICAL PHARMACOLOGY

Pharmacodynamics

Pharmacotherapeutic group

Antiepileptics, Other Antiepileptics

ATC code

N03AX14.

Mechanism of action

The active substance, levetiracetam, is a pyrrolidone derivative (S-enantiomer of α-ethyl-2-oxo- 1pyrrolidine acetamide), chemically unrelated to existing anti-epileptic active substances. The mechanism of action of levetiracetam still remains to be fully elucidated. In vitro and in vivo experiments suggest that levetiracetam does not alter basic cell characteristics and normal neurotransmission.

Levetiracetam affects intraneuronal Ca^{2+} levels by partial inhibition of N-type Ca^{2+} currents and by reducing the release of Ca^{2+} from intraneuronal stores. In addition it partially reverses the reductions in GABA- and glycine-gated currents induced by zinc and β-carbolines. Furthermore, levetiracetam has been shown to bind to a specific site in rodent brain tissue. This binding site is the synaptic vesicle protein 2A, believed to be involved in vesicle fusion and neurotransmitter exocytosis. Levetiracetam and related analogs show a rank order of affinity for binding to the synaptic vesicle protein 2A which correlates with the potency of their anti-seizure protection in the mouse audiogenic model of epilepsy. This finding suggests that the interaction between levetiracetam and the synaptic vesicle protein 2A seems to contribute to the antiepileptic mechanism of action of the medicinal product.

Pharmacodynamic effects

Levetiracetam induces seizure protection in a broad range of animal models of partial and primarily generalised seizures without having a pro-convulsant effect. The primary metabolite is inactive. In man, activity in both partial and generalised epilepsy conditions (epileptiform discharge/photoparoxysmal response) has confirmed the broad spectrum pharmacological profile of levetiracetam.

Pharmacokinetics

Levetiracetam is a highly soluble and permeable compound. The pharmacokinetic profile is linear with low intra- and inter-subject variability. There is no modification of the clearance after repeated administration. The time independent pharmacokinetic profile of levetiracetam was also confirmed following 1500 mg intravenous infusion for 4 days with twice daily dosing.

There is no evidence for any relevant gender, race or circadian variability. The pharmacokinetic profile is comparable in healthy volunteers and in patients with epilepsy.

The pharmacokinetic profile has been characterised following oral administration. A single dose of 1500 mg levetiracetam diluted in 100 ml of a compatible diluent and infused intravenously over 15 minutes is bioequivalent to 1500 mg levetiracetam oral intake, given as three 500 mg tablets. The intravenous administration of doses up to 4000 mg diluted in 100 ml of 0.9% sodium chloride infused over 15 minutes and doses up to 2500 mg diluted in 100 ml of 0.9% sodium chloride infused over 5 minutes was evaluated. The pharmacokinetic and safety profiles did not identify any safety concerns.

Adults and adolescents

Absorption

Levetiracetam is rapidly absorbed after oral administration. Oral absolute bioavailability is close to 100%. Peak plasma concentrations (C_{max}) are achieved at 1.3 hours after dosing. Steady-state is achieved after two days of a twice daily administration schedule. Peak concentrations (C_{max}) are typically 31 and 43 µg/ml following a single 1000 mg dose and repeated 1000 mg twice daily dose, respectively. The extent of absorption is dose-independent and is not altered by food.

Distribution

No tissue distribution data are available in humans. Neither levetiracetam nor its primary metabolite are significantly bound to plasma proteins (<10 %). The volume of distribution of levetiracetam is approximately 0.5 to 0.7 l/kg, a value close to the total body water volume. Peak plasma concentration (Cmax) observed in 17 subjects following a single intravenous dose of 1500 mg infused over 15 minutes was 51 ± 19 µg/mL (arithmetic average ± standard deviation).

Metabolism

Levetiracetam is not extensively metabolised in humans. The major metabolic pathway (24% of the dose) is an enzymatic hydrolysis of the acetamide group. Production of the primary metabolite, ucb L057, is not supported by liver cytochrome P450 isoforms. Hydrolysis of the acetamide group was measurable in a large number of tissues including blood cells. The metabolite ucb L057 is pharmacologically inactive.

Two minor metabolites were also identified. One was obtained by hydroxylation of the pyrrolidone ring (1.6% of the dose) and the other one by opening of the pyrrolidone ring (0.9% of the dose).

Other unidentified components accounted only for 0.6% of the dose.

No enantiomeric interconversion was evidenced in vivo for either levetiracetam or its primary metabolite.

In vitro, levetiracetam and its primary metabolite have been shown not to inhibit the major human liver cytochrome P450 isoforms (CYP3A4, 2A6, 2C9, 2C19, 2D6, 2E1 and 1A2), glucuronyl transferase (UGT1A1 and UGT1A6) and epoxide hydroxylase activities. In addition, levetiracetam does not affect the in vitro glucuronidation of valproic acid.

In human hepatocytes in culture, levetiracetam had little or no effect on CYP1A1/2, SULT1E1 or UGT1A1. Levetiracetam caused mild induction of CYP2B6 and CYP3A4. The in vitro data and in vivo interaction data on oral contraceptives, digoxin and warfarin indicate that no significant enzyme induction is expected in vivo. Therefore, the interaction of levetiracetam with other substances, or vice versa, is unlikely.

Elimination

The plasma half-life in adults was 7±1 hours and did not vary either with dose, route of administration or repeated administration. The mean total body clearance was 0.96 ml/min/kg.

The major route of excretion was via urine, accounting for a mean 95% of the dose (approximately 93% of the dose was excreted within 48 hours). Excretion via faeces accounted for only 0.3% of the dose.

The cumulative urinary excretion of levetiracetam and its primary metabolite accounted for 66% and 24% of the dose, respectively during the first 48 hours. The renal clearance of levetiracetam and ucb L057 is and

4.2 ml/min/kg respectively indicating that levetiracetam is excreted by glomerular filtration with subsequent tubular reabsorption and that the primary metabolite is also excreted by active tubular secretion in addition to glomerular filtration. Levetiracetam elimination is correlated to creatinine clearance.

Special patient populations

Children (4 to 12 years)
The pharmacokinetics in paediatric patients has not been investigated after intravenous administration. However, based on the pharmacokinetic characteristics of levetiracetam, the pharmacokinetics in adults after intravenous administration and the pharmacokinetics in children after oral administration, the exposure (AUC) of levetiracetam is expected to be similar in paediatric patients aged 4-12 years after intravenous and oral administration.

Following single dose administration (20 mg/kg) to epileptic children (6 to 12 years), the half-life of levetiracetam was 6.0 hours. The apparent body weight adjusted clearance was approximately 30% higher than in epileptic adults. Following repeated oral dose administration (20 to 60 mg/kg/day) to epileptic children (4 to 12 years), levetiracetam was rapidly absorbed. Peak plasma concentration was observed 0.5 to 1.0 hour after dosing. Linear and dose proportional increases were observed for peak plasma concentrations and area under the curve. The elimination half-life was approximately 5 hours. The apparent body clearance was 1.1 ml/min/kg.

Elderly
In the elderly, the half-life is increased by about 40% (10 to 11 hours). This is related to the decrease in renal function in this population.

Renal impairment
The apparent body clearance of both levetiracetam and of its primary metabolite is correlated to the creatinine clearance. It is therefore recommended to adjust the maintenance daily dose of levetiracetam, based on creatinine clearance in patients with moderate and severe renal impairment.

In anuric end-stage renal disease subjects the half-life was approximately 25 and 3.1 hours during interdialytic and intradialytic periods, respectively.
The fractional removal of levetiracetam was 51% during a typical 4-hour dialysis session.

Hepatic impairment
In subjects with mild (Child-Pugh A) to moderate (Child-Pugh B) hepatic impairment, the pharmacokinetics of levetiracetam were unchanged. In subjects with severe hepatic impairment (ChildPugh C), total body clearance was 50% that of normal subjects, but decreased renal clearance accounted for most of the decrease.

No dose adjustment is needed in patients with mild to moderate hepatic impairment. In patients with severe hepatic impairment, the creatinine clearance may underestimate the renal insufficiency. Therefore a 50% reduction of the daily maintenance dose is recommended when the creatinine clearance is < 60 ml/min/1.73m^2.

Gender
Levetiracetam Cmax and AUC were 20 % higher in women (N=11) compared to men (N=12). However, clearances adjusted for body weight were comparable.

Race
Formal pharmacokinetic studies of the effects of race have not been conducted. Cross study comparisons involving Caucasians (N=12) and Asians (N=12), however, show that pharmacokinetics of levetiracetam were comparable between the two races. Because levetiracetam is primarily excreted by the kidney and there are no important racial differences in creatinine clearance, pharmacokinetic differences due to race are not expected.

INCOMPATIBILITIES AND USE AND HANDLING

Intiracetam Concentrate for Solution for Infusion 100 mg/ml
See Table 1 for the recommended preparation and administration of Intiracetam Concentrate for Solution for Infusion to achieve a total daily dose of 500 mg, 1000 mg, 2000 mg or 3000 mg in two divided doses.
Table 1. Preparation and administration of Intiracetam concentrate for solution for infusion

Dose	Withdrawal Volume	Volume of Diluent	Infusion Time	Frequency of administration	Total Daily Dose

250 mg	2.5 ml (half 5 ml vial)	100 ml	15 minutes	Twice daily	500 mg/day
500 mg	5 ml (one 5 ml vial)	100 ml	15 minutes	Twice daily	1000 mg/day
1000 mg	10 ml (two 5 ml vials)	100 ml	15 minutes	Twice daily	2000 mg/day
1500 mg	15 ml (three 5 ml vials)	100 ml	15 minutes	Twice daily	3000 mg/day

This medicinal product is for single use only, any unused solution should be discarded.

Medicinal product with particulate matter or discolouration should not be used. Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

Intiracetam Concentrate for Solution for Infusion was found to be physically compatible and chemically stable when mixed with the following diluents for at least 24 hours and stored in PVC bags at controlled room temperature below 25°C.

Diluents:

- Sodium chloride 9mg/ml (0.9%) solution for injection
- Lactated Ringer's solution for injection
- Dextrose 50mg/ml (5%) solution for injection

STORAGE CONDITIONS AND EXPIRY DATE

2 years. Store below 30˚C.

From a microbiological point of view, the product should be used immediately after dilution. If not used immediately, in-use storage time and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 to 8°C.

Keep out of reach and sight of children.

Do not use after the expiry date stated on the carton box.

NATURE AND CONTENTS OF CONTAINER

Intiracetam Concentrate for Solution for Infusion is packed in glass vials with Teflon coated stoppers and sealed with an aluminium/polypropylene flip off cap. Each vial contains 5 ml of concentrate. Each carton contains 10 vials.

PRODUCT REGISTRANT

Intega Pte Ltd
100 Peck Seah Street
#08-14, PS100
Singapore 079333

Date of revision: March 2022